

Insecticide Options for Lygus Bug Management in Sugarbeets

Insecticide	Dosage in LB AI/ACRE	Product Per Acre	Restrictions on use
Asana XL <i>RUP</i>	0.03 - 0.05 lb/acre	5.8 - 9.6 fl oz	Do not apply within 21 days of harvest
Sevin XLR	1.0 - 1.5 lb/acre	2 - 3 pts	Do not apply within 28 days of harvest.
Lannate LV <i>RUP</i>	0.22 - 0.9 lb/acre	0.75 - 3.0 pts	Do not apply within 7 days of harvest. Do not feed tops to livestock within 30 days after application. Field re-entry interval is 48 hours. Fields must be posted.
Lannate SP <i>RUP</i>	0.22 - 0.9 lb/acre	0.25 - 1.0 lbs	Do not apply within 7 days of harvest. Do not feed tops to livestock within 30 days after application. Field re-entry interval is 48 hours. Fields must be posted.
Lorsban 4E	0.25 - 0.5 lb/acre	0.5 - 1.0 pts	Do not apply within 30 days of harvest. Do not apply more than 8 pints/acre (broadcast basis) per season, or make more than 4 applications per season. Do not feed tops to livestock within 30 days after application.
Malathion 57 EC	1.0 - 1.25 lb/acre	1.5 - 2.0 pts	Do not apply within 3 days of harvest

RUP - Restricted use pesticide