

2001 Insecticide Options for Tarnished Plant Bug Management in Sugarbeets.

Insecticide	Dosage in LB AI/ACRE	Product per acre	Restrictions on use
Asana XL RUP	0.03-0.05 lb/acre	5.8-9.6 fl oz	Do not apply within 21 days of harvest
Sevin XLR	1.0-1.5 lb/acre	2-3 pts	Do not apply within 28 days of harvest
Lannate LV RUP	0.22-0.9 lb/acre	0.75-3.0 pts	Do not apply within 7 days of harvest. 0.25-0.5 lb/acre Field re-entry intervals is 48 hrs. Fields must be posted.
Lannate SP RUP	0.22-0.9 lb/acre	0.25-0.5 lbs	Do not apply within 7 days of harvest. Do not feed tops to livestock within 30 days after application. Field re-entry intervals is 48 hrs. Fields must be posted
Lorsban 4E	0.25-0.5 lb/acre	0.5-1.0 pts	Do not apply within 30 days of harvest. Do not apply more than 8 pints/acre (broadcast basis) per season, or make more that 4 applications per season. Do not feed tops to livestock within 30 days after application.
Malathion 57EC	1.0-1.25 lb/acre	1.5-2.0 pts.	Do not apply within 3 days of harvest.

RUP - Restricted use pesticide